arXiv:0801.3733v1 [astro-ph] 24 Jan 2008

Consistent simulations of substellar atmospheres and non-equilibrium dust-cloud formation

Christiane Helling

SUPA, School of Physics & Astronomy, Univ. of St Andrews, North Haugh, St Andrews, KY16 9SS, UK

Christiane.Helling@st-and.ac.uk

Matthias Dehn

Hamburger Sternwarte, Gojenbergsweg 112, 21029 Hamburg, Germany

Peter Woitke

UK ATC, Royal Observatory, Blackford Hill, Edinburgh EH9 3HJ, UK

and

Peter H. Hauschildt

Hamburger Sternwarte, Gojenbergsweg 112, 21029 Hamburg, Germany

ABSTRACT

We aim to understand cloud formation in substellar objects. We combined the non-equilibrium, stationary cloud model of Helling, Woitke & Thi (2008; seed formation, growth, evaporation, gravitational settling, element conservation) with the general-purpose model atmosphere code PHOENIX (radiative transfer, hydrostatic equilibrium, mixing length theory, chemical equilibrium) in order to consistently calculate cloud formation and radiative transfer with their feedback on convection and gas phase depletion. We calculate the complete 1D model atmosphere structure and the chemical details of the cloud layers. The DRIFT-PHOENIX models enable the first stellar atmosphere simulation that is based on the actual cloud *formation* process. The resulting (T, p) profiles differ considerably from the previous limiting PHOENIX cases DUSTY and COND. A tentative comparison with observations demonstrates that the determination of effective temperatures based on simple cloud models has to be applied with care. Based on our new models, we suggest a mean $T_{\rm eff} = 1800$K for the L - dwarf twin-binary system DENIS J0205-1159 which is up to 500K hotter than suggested in the literature. We show transition spectra for gas-giant planets which form dust clouds in their atmospheres and evaluate photometric fluxes for a WASP-1 type system.

Subject headings: astrochemistry — stars: atmospheres — methods: numerical

1. Introduction

Today's most efficient tools to interpret the observed spectra of substellar objects, i.e., brown dwarfs and planets, are 1D static atmosphere simulations. Comparisons with observations, based on the solution of the radiative-transfer problem which is done in great detail with respect to the gas-phase opacities (Tsuji 2002, 2005; Allard et al. 2001, Ackerman & Marley 2001, Burrows & Sharp 1999, Barman et al. 2005), ideally yield insight into the atmospheric structure and chemistry providing finger prints of its evolutionary state.

Brown dwarf and planetary atmospheres have a far more complex chemistry than stellar objects due to the formation of clouds, which bind chemical elements and, hence, strongly influences the remnant gas phase inside the atmosphere. The presence of such clouds was previously simplified in static model atmosphere codes. We now move a significant step forward by kinetically treating the chemistry of cloud formation as a phase-non-equilibrium process in the framework of model atmosphere simulations. Our dust model has been studied so far for a given $(T, p, v_{\rm conv})$ structure (T - gas temperature, p - gas pressure, $v_{\rm conv}$ - convective velocity), and we present in this letter the first consistent simulation of cloud micro-physics and atmospheric structure which allows for the first time to study the feedback of the dust *formation* onto the atmospheric structure. We present our first consistent DRIFT-PHOENIX results together with a tentative comparison to the observed spectrum of DENIS J0205–1159 and point out possible uncertainties in present $T_{\rm eff}$ determinations (Sect. 4). We, however, leave aside the issue of hydrodynamical cloud formation (e.g. Showman et al. 2006, Knutson et al. 2007, Rauscher et al. 2007) which unavoidably has to deal with the turbulent closure problem (Helling et al. 2004, Helling 2007). We calculate transition spectra for gas-giant planets and a WASP-1 type star, and evaluate the photometric fluxes for the 2MASS system, for the VISIR system, and for the IRAC band systems.

2. Method

We combined our model describing non-equilibrium dust formation (DRIFT; Woitke & Helling 2003, 2004; Helling & Woitke 2006; Helling, Woitke, Thi 2008) with our general-purpose model atmosphere code (PHOENIX; Hauschildt & Baron 1999). We consider a stationary dust formation process, where seeds form (homo-molecular TiO_2-nucleation, see Woitke & Helling 2004, Sect. 2.2) from a highly supersaturated gas, grow to macroscopic

particles of μm size, gravitationally settle into deeper layers, and eventually evaporate as the local temperature becomes too high for thermal stability (Woitke & Helling 2003). Coagulation, a process amongst existing grains, is omitted so far since it acts on longer time and size scales than the dust formation processes considered here. Typical time scales to form a 0.1μm SiO_2-grain in a gas of $T = 1500$K, $\rho = 10^{-6}$ g cm^{-3} by continuous growth and coagulation are $\tau_{\rm gr} \approx 10^{-1}$s (Fig. 3, Woitke & Helling 2003) and $\tau_{\rm coag} \approx 10$ s (Eq. 37[1], Rossow 1978), respectively. The settling time scale is $\tau_{\rm grav} \approx 10^5$s (Eq. 20, Woitke & Helling 2003) for this grain crossing a pressure scale hight ($H_{\rm p} = 10^6$cm) assuming a constant drift velocity $v_{\rm drift} \approx 10$cm s^{-1}. $v_{\rm drift}(a)$ rapidly increases for $\rho > 10^{-6}$ g cm^{-3}, hence $\tau_{\rm grav}$ decreases inward.

A truly static atmosphere would not contain any dust (Woitke & Helling 2004), therefore we include mixing by convection and overshooting by assuming an exponential decrease of the mass exchange frequency in the radiative zone (Eq. 9 in (Woitke & Helling 2004) with $\beta = 2.2$ and $\tau_{\rm mix}^{\rm min} = 2/(H_{\rm p} v_{\rm conv})$), which serves to replenish the upper atmospheres and keeps the cycle of dust formation running. In order to keep the computing time reasonable, we consider the growth/evaporation of 7 solids ($MgSiO_3$[s], Mg_2SiO_4[s], MgO[s], SiO_2[s], SiO[s], Al_2O_3[s], TiO_2[s]) made of 5 different elements for which we solve 15 possibly stiff conservation equations for 256 atmospheric layers. The whole 1D atmosphere problem is solved iteratively in DRIFT-PHOENIX where PHOENIX provides the actual $(T, p, v_{\rm conv})$ structure and DRIFT solves the dust moment and element conservation equations in the subsonic, large Knudsen number case. DRIFT hands back $(f(V,z), V_{\rm s}(z), \epsilon_{\rm i(i=Mg,Si,O,Al,Ti)})$ with $f(V,z)$ the parameterised distribution function of grain volume V at atmospheric height z, $V_{\rm s}$ the chemical dust composition in volume fractions for all solids s involved, and $\epsilon_{\rm i}$ the remaining element abundances for all involved elements i in the gas phase (Dehn 2007). Effective medium and Mie theory are used to calculate the opacity of these composite and chemically heterogeneous dust grains in PHOENIX (see Helling, Woitke & Thi (2008), Table 2).

3. Results

Figure 1 shows that DRIFT-PHOENIX (T, p)-profiles for late L-type brown dwarfs basically fall between the two limiting cases of the previous PHOENIX versions DUSTY[2] and COND[3] (Allard et al. 2001). DUSTY-PHOENIX models are hotter and COND-PHOENIX models are cooler than our self-consistent model. We notice that, amongst the ($T_{\rm eff} = 1800$K,

[1] All input quantities are taken from the model presented in this paper.

[2] DUSTY: Dust is considered as opacity source and element sink.

[3] COND: Dust is considered as element sink only.

log g = 5.0) models, the DRIFT-PHOENIX profile has the highest temperatures inside the convectively active region below 1 bar. For a given local temperature, the atmospheric pressure increases inside the dust forming region at $T > 1000K$ with increasing surface gravity as suggested by previous models. The result is a cloud being thermally stable at higher temperatures, hence sitting deeper in the atmosphere.

Grain sizes $\langle a \rangle$, material composition $V_\mathrm{s}/V_\mathrm{tot}$, number of dust particles n_d and remaining element abundances ϵ_i determine the gas-phase opacity and are major ingredients for the radiative transfer calculation. Figure 2 shows for the model with ($\mathrm{T_{eff}}$ = 1800K, log g = 5.0) that the upper cloud layer is mainly made of small silicate grains with only some MgO[s] and Fe[s] impurities which is in accordance with our previous results (Helling et al. 2006, Helling, Woitke & Thi 2008). However, only a few of such silicate grains populate the cloud deck (Fig. 3 thick gray line). The material composition changes drastically at $\sim$ 0.1bar where big grains form the majority of the grain population. The dust cloud formation causes a local element sink (Fig. 3). The strongest element depletion coincides with the maximum number of particles in pressure space. High-temperature condensable elements like Al and Ti are depleted in a wider atmosphere range since the associated solids are thermally stable over a larger temperature range than those binding Mg and Si.

4. Application

Brown Dwarf spectra: DRIFT-PHOENIX was used to tentatively reproduce the observed spectrum of the L - dwarf twin-binary system DENIS J0205-1159 observed between 0.5μm and 2.6 μm (Reid et al. 2001; Leggett et al. 2001). The determination of its mean effective temperatures proves difficult: Leggett et al. (2001) suggested $\mathrm{T_{eff}}$ =1900K and log g = 5.5 based on DUSTY-PHOENIX models and $\mathrm{T_{eff}}$ = 1400 ... 1600K from structural models. Golimowsky et al. (2004) suggest, depending on the system's age, an effective temperature between 1350K and 1700K. Vrba et al. (2004) suggest $\mathrm{T_{eff}}$ =1563K with an astonishing precision. Brandeker et al. (2006) report similar difficulties in determining stellar parameter of the binary OPH 162225-240515. Based on DUSTY-PHOENIX and Burrows models (Burrow et al. 2006) they suggest $\mathrm{T_{eff}}$ = (2350$\pm$150)K and $\mathrm{T_{eff}}$ = (2100$\pm$100)K for their components A an B, respectively.

Figure 4 shows our comparison of DRIFT-PHOENIX synthetic spectra of $\mathrm{T_{eff}}$ =1800K with log g = 5.0 / log g = 6.0 (red solid / green dash-dot line) and $\mathrm{T_{eff}}$ =1900K with log g = 5.5 (blue dotted) to the DENIS J0205-1159 spectrum (gray) between 0.5μm and 2.6 μm. The synthetic spectra are normalised to the observed spectrum in the K-band at

$\lambda = 2.1\,\mu$m ([4]). The stellar parameters have been chosen to produce the overall best fit in this spectral range but we cannot identify a single model that fits the whole wavelength interval. The J-band data are best reproduced by the (T_{eff} =1800K, log $g = 6.0$) model. The H- and K-band data are best reproduced by the (T_{eff} =1800K, log $g = 5.0$) model. Note that the H band is most sensitive to chances in the surface gravity of the models. The (T_{eff} =1900K, log $g = 5.5$) model in general produces too much flux except for the long-wavelength edge of the K band. Tsuji (2005) and Cushing et al. (2007) report similar challenges for their models in simultaneous fitting the JHK bands. All shown models reproduce very well the H_2O absorption edge longword 1.3 μm and 1.75 μm.

We tentatively suggest an effective temperature of T_{eff} =1800K. Our tentative suggestion for the surface gravity is log $g = 5.0$ since an increase in log g broadens the H band peak and a decrease would narrow it, both causing even stronger discrepancies to the observed spectrum. It appears unlikely, according to our model, that the system's mean temperature should be as low as $T_{eff} \lesssim 1600$K, since the synthetic flux would drop considerably.
Our results furthermore suggest that L - dwarfs contain a considerable amount of dust in their atmospheres with a gradually changing chemical dust composition (see Fig. 2). This explains also why our simulations yield about the same T_{eff} range as those in Leggett et al. (2001) who applied the DUSTY-PHOENIX limiting case, where the dust was kept in the atmosphere as opacity source, and consequently a high atmospheric dust content was assumed. The actual (T, p)-profiles, however, are very different with all its implications for the gas-phase and cloud formation chemistry (see Fig. 1). Compared to $T_{eff} = 1350 - 1700$K (Golimowsky et al. 2004), our results would then suggest an age of 2 Gyr.
So far, our synthetic spectra are the results of a detailed micro-physical treatment of dust formation in phase-non-equilibrium including the formation of seed particles, their growth, evaporation, and the effect of gravitational settling. Grain sizes and grain size distribution, cloud thicknesses, dust composition, total dust content etc. are results of the simulations and are not prescribed as it was needed in the DUSTY- and COND-PHOENIX models. Our simulations, however, leave space for further improvement of our synthetic spectra like for instance regarding the treatment of alkali line profiles in the optical (Allard et al. 2007, Johnas et al. 2008), the formation of liquid water droplets, or the improvement of our mixing modelling.

Gas-giant transition spectra: We have calculated transition spectra for a hot and a medium temperature gas-giant planet with $T_{eff} = 1800$K and $T_{eff} = 1400$K, respectively,

[4]Cushing et al. (2007) normalise to 1.25μm in the J, 1.6μm in the H, and 2.1μm in the K band.

and $\log g = 3.0$ in the light of a WASP-1 type host star ($T_{\rm eff} = 6140$K, $\log g = 4.31$, solar metallicity; compare Collier Cameron et al. 2007). We neglect, as a first attempt, the irradiation of the host star and day-night effects. For example, Barman et al. (2005) and Burrows et al. (2006) include these effects but do not account for the influence of cloud formation. Figure 5 shows $D \times F_\lambda^{\rm planet}/F_\lambda^{\rm star}$, the transition spectra between $0.5\,\mu$m and $15\,\mu$m corrected by the surface area ratio $D = (R_{\rm planet}/R_{\rm star})^2$ ($R_{\rm planet} = 8\,10^9$cm$= 1.11\,R_{\rm Jupiter}$, $R_{\rm star} = 8.63\,10^{10}cm= 1.24\,R_\odot$; $M_{\rm planet} = 0.5\,M_{\rm Jupiter}$, F_λ - surface flux). We add $D \times F_{\rm bb}(T_{\rm eff}^{\rm planet})/F_{\rm bb}(T_{\rm eff}^{\rm star})$ and the photometric fluxes for the planets thermal emission for the VISIR and IRAC bands, and the 2MASS J H Ks bands for comparison. The transit contrast due to the thermal emission increases with increasing wavelength but the difference between the $T_{\rm eff} = 1800$K and $T_{\rm eff} = 1400$K model become constant for $\lambda > 8\mu$m. Our cloud-covered gas-giant planets produce photometric contrasts which are comparable to the secondary eclipse measurements by Charbonneau et al. (2005), Deming et al. (2007), Snellen & Covino (2007). We furthermore note a good agreement with the results on HD 209458b by Barman et al. (2005) despite all the uncertainties in the cloud and the day-night-effect modelling.

5. Summary

The modelling of cloud formation in substellar atmospheres has a strong impact on the objects temperature-pressure structure which in turn determines the cloud's chemical composition, the grain size distribution function, and the cloud's location inside the atmosphere. These are results of our consistent simulation of substellar atmospheres and detailed non-equilibrium dust cloud formation. We demonstrate synthetic transition spectra for gas-giant planets and calculate synthetic photometric fluxes based on our consistent solution of dust cloud formation and radiative transfer problem. A future goal is to extend our model to solar-system-like planets with much cooler atmosphere.

As we have shown for the field object Denis J0205-1159, stellar parameter determinations based on a comparison with synthetic spectra can vary considerable. As one consequence of this, we have set out to conduct a component-based study [5] where e.g. cloud compositions, dust-to-gas-ratios, and grain sizes are compared for different cloud models (Helling et al. 2007, 2008).

We thank Aleks Scholz, Andrew Collier Cameron and the anonymous referee for helpful

[5] http://phoenix.hs.uni-hamburg.de/BrownDwarfsToPlanets1/

discussions on the paper's subject. The computer support at the School of Physics and Astronomy St Andrews is highly acknowledged. Most of the literature search was done with ADS.

REFERENCES

Allard F., Hauschildt P.H., Alexander D.R. et al. 2001, ApJ 556, 357

Allard F., Allard N., Homeier D. et al. 2007, A&A, in press

Ackerman A., Marley M., 2001, ApJ 556, 872

Barman T., Hauschildt P.H., Allard F. 2005 ApJ 632, 1132

Brandeker A. Jayawardhana R. Ivanov V.D, Kurtev R. 2006, ApJ653, L61

Burrows A., Sharp C.M. 1999, ApJ 512, 843

Burrows A., Sudarsky D., Hubeny I. 2006, ApJ 650, 1140

Charbonneau D., Allen L.E., Megeath S.T. et al. 2005, ApJ 626, 523

Collier Cameron A., Bouchy F., Hébrad G. et al. 2007, MNRAS 375, 951

Cushing M.C., Marley M.S., Saumon D et al. 2007, ApJ, astro-ph arXiv:0711.0801v1

Dehn M. 2007, PhD Thesis, University Hamburg

Deming D., Richardson L.J., Harrington J. 2007, MNRAS 378, 148

Hauschild P., Baron E. 1999, JCAM 109, 41

Helling Ch., Klein R., Woitke P., Nowak U., Sedlmayr E. 2004, A&A 423, 657

Helling Ch., Woitke, P. 2006, A&A 455, 325

Helling Ch. 2007, in *Conv. in Astrophys.*, eds. Kupka, Roxburgh, Chan, 224

Helling Ch., Woitke P., Thi W.-F. 2008, A&A submitted

Helling Ch., Ackerman A., Allard F., Dehn M., Hauschildt P., Homeier D., Lodders K., Marley M., Rietmeijer F., Tsuji T, Woitke P. 2007, arXiv:0711.3993

Helling Ch., Ackerman A., Allard F., Dehn M., Hauschildt P., Homeier D., Lodders K., Marley M., Rietmeijer F., Tsuji T, Woitke P. 2008, A&A, in preperation

Johnas C.M.S., Helling, Ch., Dehn, M., Woitke, P. Hauschildt P.H. 2008, MNRAS Letter, in press (arXiv:0801.3544)

Knutson H.A., Charbonneau D., Allen L.E., et al. 2007, Nature 477 (7141), 183

Leggett S.K., Allard F., Geballe T.R. et al. 2001, ApJ 548, 908

Rauscher E., Menou K., Cho J.Y.-K., Seager S., Hansen B.M.S. 2007, ApJ 662(2), L115

Reid I.N., Burgasser A.J., Cruz K.L., Kirkpatrick J.D., Gizis J.E. 2001, ApJ 121, 1710

Rossow, W.B. 1978, Icarus 36, 1

Showman A.P., Gierasch P.J., Lion Y. 2006, Icarus 182(2), 513

Snellen I.A.G., Covino E. 2007, MNRAS 375, 307

Tsuji T. 2005, ApJ 621, 1033

Woitke P., Helling Ch. 2003, A&A 399, 297

Woitke P., Helling Ch. 2004, A&A 414, 335

This preprint was prepared with the AAS LaTeX macros v5.2.




Fig. 1.— Temperature-pressure (T, p) profiles for $T_{\rm eff}$ = 1800K, log g = 5.0 (solar abundances) as result of different dust modelling (DRFT, DUSTY, COND) in the same model atmosphere code (PHOENIX). We show for comparison models with ($T_{\rm eff}$ = 1800K, log g = 6.0) and ($T_{\rm eff}$ = 1900K, log g = 5.5); see also Fig. 4.




Fig. 2.— Material composition of the dust cloud in volume fractions $V_{\rm s}/V_{\rm tot}$ ($V_{\rm s}$ - volume fraction of solid s, $V_{\rm tot}$ - total dust volume) and the mean particle size $\langle a \rangle$ [μm] (thick gray) for the ($T_{\rm eff}$ = 1800K, log g = 5.0) model with solar element abundances.




Fig. 3.— Metalicity $\log(\epsilon_{\rm i}^0/\epsilon_{\rm H}) - \log(\epsilon_{\rm i}/\epsilon_{\rm H})$ ($\epsilon_{\rm i}^0$ - solar abundances of element i, H - hydrogen) and dust number density $n_{\rm d}$ [cm^{-3}] (thick gray) for the ($T_{\rm eff} = 1800$K, $\log g = 5.0$) model.




Fig. 4.— Drift-Phoenix synthetic spectra in comparison to the brown dwarf twin-binary Denis J0205-1159 spectrum from Reid et al. (2001). The synthetic spectra are normalised to the observed spectrum at $\lambda = 2.1\mu$m, they are Gaussian convolved to 3400 sampling points with a spectral resolution of R=600. For (T, p) structures see Fig. 1.




Fig. 5.— Transition spectra for two gas-giant planets seen in the light of a WASP-1 type host star (T_{eff} = 6140K, log g=4.31; compare Collier Cameron et al. 2007). The synthetic spectra are Gauss convolved to 3000 sampling points with R=15000. Horizontal lines indicate the photometric bands and the symbols give photometric fluxes for the J, H, K bands (■), the VISIR bands (⧫) PAH1, ArIII (1), SIV (2), PAH2, SiV (3), SiC, and the IRAC bands Band 1 – 4 (▲).